

BOUNDARIES OF REDUCED FREE GROUP C^* -ALGEBRAS

NARUTAKA OZAWA

ABSTRACT. We prove that the crossed product C^* -algebra $C_r^*(\Gamma, \partial\Gamma)$ of a free group Γ with its boundary $\partial\Gamma$ naturally sits between the reduced group C^* -algebra $C_r^*\Gamma$ and its injective envelope $I(C_r^*\Gamma)$. In other words, we have natural inclusion $C_r^*\Gamma \subset C_r^*(\Gamma, \partial\Gamma) \subset I(C_r^*\Gamma)$ of C^* -algebras.

1. INTRODUCTION

A deep theorem of Kirchberg [Ki][KP] states that for any separable exact C^* -algebra B , there exists a separable nuclear C^* -algebra A that contains B as a C^* -subalgebra. (Indeed, one can take A as the Cuntz algebra $\mathcal{O}_2$.) In this note, we are interested in finding a more tight (or canonical) embedding of a given separable exact C^* -algebra into a separable nuclear C^* -algebra. This is partly motivated by Hamana's theory of injective envelopes. An embedding of B into A is called rigid if the identity map on A is the only completely positive map on A which restricts to the identity map on B . Hamana [Ha] proved that for any C^* -algebra B , there exists an injective C^* -algebra $I(B)$, called the injective envelope of B , with a rigid embedding of B into $I(B)$. We note that the injective envelope $I(B)$ is unique up to natural identification.

Let Γ be the free group of rank $2 \leq r < \infty$ and $\partial\Gamma$ be its boundary, i.e., the compact topological space of all one-sided infinite reduced words, equipped with the product topology. The free group Γ acts on $\partial\Gamma$ by translation and this action is topologically amenable (cf. [AR]). The reduced group C^* -algebra $C_r^*\Gamma$ is naturally contained in the reduced crossed product C^* -algebra $C_r^*(\Gamma, \partial\Gamma)$, which is nuclear. A measure μ on $\partial\Gamma$ is called *quasi-invariant* if for any measurable subset $X \subset \partial\Gamma$ and any $s \in \Gamma$, one has $\mu(sX) = 0$ if and only if $\mu(X) = 0$. A measure μ on $\partial\Gamma$ is called *doubly-ergodic* if the diagonal action of Γ on $(\partial\Gamma^2, \mu^{\otimes 2})$ is ergodic. Fix a quasi-invariant and doubly-ergodic measure μ on $\partial\Gamma$. For instance, the harmonic measure of any non-degenerate symmetric random walk on Γ has this property (Theorem 2.8 in [Ka]). Associated with μ , there is a natural inclusion of $C_r^*(\Gamma, \partial\Gamma)$

Date: November 20, 2004.

2000 Mathematics Subject Classification. Primary 46L05; Secondary 20F67.

Key words and phrases. free groups, reduced group C^* -algebras, injective envelopes.

The author was supported in part by JSPS.

into the crossed product von Neumann algebra $\Gamma \ltimes L^\infty(\partial\Gamma, \mu)$, which is injective;

$$C_r^*\Gamma \subset C_r^*(\Gamma, \partial\Gamma) \subset \Gamma \ltimes L^\infty(\partial\Gamma, \mu).$$

See [Ok] [RR] for information on the type of the factor $\Gamma \ltimes L^\infty(\partial\Gamma, \mu)$.

We state the main theorem of this note. A somewhat similar result was proved in [Sp].

Theorem 1. *Let Γ be the free group of rank $2 \leq r < \infty$, $\partial\Gamma$ be its boundary and μ be a quasi-invariant and doubly-ergodic measure on $\partial\Gamma$. If*

$$\theta: C_r^*(\Gamma, \partial\Gamma) \rightarrow \Gamma \ltimes L^\infty(\partial\Gamma, \mu)$$

is a completely positive map with $\theta|_{C_r^\Gamma} = \text{id}_{C_r^*\Gamma}$, then $\theta = \text{id}$.*

Corollary 2. *Let Γ and $\partial\Gamma$ be as above. Then, the nuclear C^* -algebra $C_r^*(\Gamma, \partial\Gamma)$ sits (as a C^* -algebra) between $C_r^*\Gamma$ and its injective envelope $I(C_r^*\Gamma)$, i.e., we have natural inclusions $C_r^*\Gamma \subset C_r^*(\Gamma, \partial\Gamma) \subset I(C_r^*\Gamma)$ of C^* -algebras.*

The author conjectures that for any separable exact C^* -algebra B , there exists a nuclear C^* -algebra between B and its injective envelope $I(B)$.

The theorem is a consequence of the following proposition from ergodic theory, inspired by [MS]. We recall that a map between spaces with Γ -actions is called Γ -equivariant if it commutes with the Γ -actions.

Proposition 3. *Let Γ be the free group of rank $2 \leq r < \infty$, $\partial\Gamma$ be its boundary and μ be a quasi-invariant and doubly-ergodic measure on $\partial\Gamma$. If*

$$\varphi: C(\partial\Gamma) \rightarrow L^\infty(\partial\Gamma, \mu)$$

is a unital positive Γ -equivariant map, then $\varphi = \text{id}$.

Actually the same result *mutatis mutandis* holds for a discrete subgroup Γ of the group of isometries on an exponentially-bounded Gromov hyperbolic space $\mathcal{X}$ and a closed subspace of $\partial\mathcal{X}$ on which Γ acts minimally. It is plausible that the assumption on μ can be relaxed to that μ is quasi-invariant and diffuse. (For this matter, it suffices to show this assertion under the additional assumption that φ is a Γ -equivariant $*$ -homomorphism.)

2. PROOF OF PROPOSITION

Let Γ be the free group of rank $2 \leq r < \infty$, $\partial\Gamma$ be its boundary and μ be a quasi-invariant and doubly-ergodic measure on $\partial\Gamma$. Let a unital positive Γ -equivariant map $\varphi: C(\partial\Gamma) \rightarrow L^\infty(\partial\Gamma, \mu)$ be given.

We denote by $\mathcal{M}(\partial\Gamma)$ the space of all finite Borel measures on $\partial\Gamma$ and let $\mathcal{M}_{\leq 2}(\partial\Gamma)$ be its subset of measures which are supported on at most two points. It is well-known (cf. Section 5 in [Ad]) that $\mathcal{M}_{\geq 3}(\partial\Gamma) = \mathcal{M}(\partial\Gamma) \setminus \mathcal{M}_{\leq 2}(\partial\Gamma)$ is a tame Γ -space. From this and the double ergodicity of μ , it follows that any Γ -equivariant Borel map from $\partial\Gamma^2$ into $\mathcal{M}_{\geq 3}(\partial\Gamma)$ has image in a single orbit $\mu^{\otimes 2}$ -a.e.

Since every non-trivial element of Γ acts hyperbolically on $\partial\Gamma$, the stabilizer of this orbit is trivial. Since $\mu^{\otimes 2}$ is diffuse, the pre-image of any point of this orbit can be divided into disjoint non-null pieces, contradicting ergodicity of $\mu^{\otimes 2}$. Thus any Γ -equivariant Borel map from $\partial\Gamma^2$ into $\mathcal{M}(\partial\Gamma)$ has image in $\mathcal{M}_{\leq 2}(\partial\Gamma)$ $\mu^{\otimes 2}$ -a.e.

Fix a dense Γ -invariant $*$ -subalgebra $\mathcal{C}$ in $C(\partial\Gamma)$ which is algebraically (over $\mathbb{C}$) generated by a countable set. Then, there exists a Γ -equivariant Borel map

$$\varphi_*: \partial\Gamma \ni \xi \mapsto \varphi_*^\xi \in \mathcal{M}(\partial\Gamma)$$

such that for μ -a.e. $\xi \in \partial\Gamma$, we have

$$\forall f \in \mathcal{C} \quad \int f(\eta) d\varphi_*^\xi(\eta) = \varphi(f)(\xi).$$

We consider the Γ -equivariant Borel map

$$\partial\Gamma^2 \ni (\xi, \eta) \mapsto \varphi_*^\xi + \delta_\xi + \delta_\eta \in \mathcal{M}(\partial\Gamma),$$

where δ_ξ is the Dirac measure on $\xi \in \partial\Gamma$. By the previous result, this map has image in $\mathcal{M}_{\leq 2}(\partial\Gamma)$ $\mu^{\otimes 2}$ -a.e. Since the diagonal subset of $\partial\Gamma^2$ is $\mu^{\otimes 2}$ -null and ξ and η are independent, we conclude that $\varphi_*^\xi = \delta_\xi$ for μ -a.e. $\xi \in \partial\Gamma$. This means that $\varphi = \text{id}$. $\square$

3. PROOF OF THEOREM

Let Γ be the free group of rank $2 \leq r < \infty$, $\partial\Gamma$ be its boundary and μ be a quasi-invariant and doubly-ergodic measure on $\partial\Gamma$. For $s \in \Gamma$ and $f \in L^\infty(\partial\Gamma, \mu)$, we define $s \cdot f \in L^\infty(\partial\Gamma, \mu)$ by $(s \cdot f)(\xi) = f(s^{-1}\xi)$. The crossed product von Neumann algebra $\Gamma \ltimes L^\infty(\partial\Gamma, \mu)$ is generated (as a von Neumann algebra) by a unitary representation λ of Γ and $L^\infty(\partial\Gamma, \mu)$. They satisfy the covariance property that $\lambda(s)f\lambda(s)^* = s \cdot f$ for $s \in \Gamma$ and $f \in L^\infty(\partial\Gamma, \mu)$. Moreover, there exists a normal faithful conditional expectation E from $\Gamma \ltimes L^\infty(\partial\Gamma, \mu)$ onto $L^\infty(\partial\Gamma, \mu)$ such that for $s \in \Gamma$ and $f \in L^\infty(\partial\Gamma, \mu)$, one has $E(\lambda(s)f) = f$ if $s = 1$ and $E(\lambda(s)f) = 0$ if $s \neq 1$. The C^* -subalgebra in $\Gamma \ltimes L^\infty(\partial\Gamma, \mu)$ generated by $\lambda(\Gamma)$ and $C(\partial\Gamma)$ is naturally $*$ -isomorphic to the reduced crossed product C^* -algebra $C_r^*(\Gamma, \partial\Gamma)$ and the C^* -subalgebra in $C_r^*(\Gamma, \partial\Gamma)$ generated by $\lambda(\Gamma)$ is naturally $*$ -isomorphic to the reduced group C^* -algebra $C_r^*\Gamma$;

$$C_r^*\Gamma \subset C_r^*(\Gamma, \partial\Gamma) \subset \Gamma \ltimes L^\infty(\partial\Gamma, \mu).$$

We prove Theorem 1 and Corollary 2.

Proof of Theorem 1. Let $\theta: C_r^*(\Gamma, \partial\Gamma) \rightarrow \Gamma \ltimes L^\infty(\partial\Gamma, \mu)$ be a completely positive map such that $\theta|_{C_r^*\Gamma} = \text{id}_{C_r^*\Gamma}$. Consider the unital positive map $\varphi = E \circ \theta|_{C(\partial\Gamma)}$.

Since $\theta|_{C_r^*\Gamma} = \text{id}_{C_r^*\Gamma}$, the completely positive map θ is automatically a $C_r^*\Gamma$ -bimodule map (cf. [CE]). Hence, for any $s \in \Gamma$ and $f \in C(\partial\Gamma)$, we have

$$\begin{aligned}\varphi(s \cdot f) &= E(\theta(\lambda(s)f\lambda(s)^*)) \\ &= E(\lambda(s)\theta(f)\lambda(s)^*) \\ &= s \cdot \varphi(f),\end{aligned}$$

where we note that $E(\lambda(s)x\lambda(s)^*) = s \cdot E(x)$ for any $x \in \Gamma \ltimes L^\infty(\partial\Gamma, \mu)$. It follows from Proposition 3 that $\varphi = \text{id}$. Since E is faithful, this implies that $\theta|_{C(\partial\Gamma)} = \text{id}_{C(\partial\Gamma)}$ and the conclusion follows. $\square$

Proof of Corollary 2. Since $\Gamma \ltimes L^\infty(\partial\Gamma, \mu)$ is injective, the injective envelope $I(C_r^*\Gamma)$ sits (as an operator system) between $C_r^*\Gamma$ and $\Gamma \ltimes L^\infty(\partial\Gamma, \mu)$. By Theorem 1, a completely positive projection θ from $\Gamma \ltimes L^\infty(\partial\Gamma, \mu)$ onto $I(C_r^*\Gamma)$ acts identically on $C_r^*(\Gamma, \partial\Gamma)$. It follows that $C_r^*(\Gamma, \partial\Gamma)$ is contained in $I(C_r^*\Gamma)$. This inclusion agrees with each C^* -algebra structure, because the C^* -algebra structure of $I(C_r^*\Gamma)$ comes from the Choi-Effros product [CE] associated with θ . $\square$

REFERENCES

- [Ad] S. Adams, *Reduction of cocycles with hyperbolic targets*. Ergodic Theory Dynam. Systems **16** (1996), 1111–1145.
- [AR] C. Anantharaman-Delaroche and J. Renault. *Amenable groupoids*. With a foreword by Georges Skandalis and Appendix B by E. Germain. Monographies de L'Enseignement Mathématique **36**. Geneva, 2000.
- [CE] M.-D. Choi and E. G. Effros, *Injectivity and operator spaces*. J. Funct. Anal. **24** (1977), 156–209.
- [Ha] M. Hamana, *Injective envelopes of C^* -algebras*. J. Math. Soc. Japan **31** (1979), 181–197.
- [Ka] V. A. Kaimanovich, *Ergodicity of harmonic invariant measures for the geodesic flow on hyperbolic spaces*. J. Reine Angew. Math. **455** (1994), 57–103.
- [Ki] E. Kirchberg, *Exact C^* -algebras, tensor products, and the classification of purely infinite algebras*. Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994), 943–954, Birkhäuser, Basel, 1995.
- [KP] E. Kirchberg and N. C. Phillips, *Embedding of exact C^* -algebras in the Cuntz algebra $\mathcal{O}_2$* . J. Reine Angew. Math. **525** (2000), 17–53.
- [MS] N. Monod and Y. Shalom, *Cocycle superrigidity and bounded cohomology for negatively curved spaces*. J. Differential Geom. **67** (2004), 395–455.
- [Ok] R. Okayasu, *Type III factors arising from Cuntz-Krieger algebras*. Proc. Amer. Math. Soc. **131** (2003), 2145–2153.
- [RR] J. Ramagge and G. Robertson, *Factors from trees*. Proc. Amer. Math. Soc. **125** (1997), 2051–2055.
- [Sp] J. Spielberg, *Free-product groups, Cuntz-Krieger algebras, and covariant maps*. Internat. J. Math. **2** (1991), 457–476.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, KOMABA, 153-8914,
DEPARTMENT OF MATHEMATICS, UCLA, LOS ANGELES, CA 90095-1555
E-mail address: narutaka@ms.u-tokyo.ac.jp